

FANO VARIETIES WITH LARGE DEGREE ENDOMORPHISMS

JÁNOS KOLLÁR AND CHENYANG XU

We work over the complex number field $\mathbb{C}$. An *endomorphism* of a projective variety X is a *morphism* $f : X \rightarrow X$. If f is surjective, then it is automatically finite.

Examples of projective varieties with an endomorphism of degree > 1 are $\mathbb{P}^n$ and abelian varieties. In fact, these are essentially all known smooth examples. When X has non-negative Kodaira dimension, this is discussed in [NZ07]. It is also conjectured that if X is a smooth Fano variety of Picard number 1 that admits a degree > 1 endomorphism, then $X \cong \mathbb{P}^n$ [Am97]. This is confirmed in some cases, including when the dimension is less or equal to 3 [ARV99], when X is a quasi-homogeneous [HN08], when X is a large degree hypersurface of a prime Fano manifold of Picard number 1 [Ch08], or when X contains a smooth conic [HM03]. [Zh08] considered Fano varieties with Picard number 1 and, with terminal singularities that have an endomorphism of degree > 1 . He conjectured that they are necessarily rational. Our aim in this note is to point out that this is not quite true and suggest a slight modification that is consistent with these examples.

1. **Example.** Any subgroup $G \subset S_{n+1}$ acts on $\mathbb{P}^n$ where S_{n+1} acts by permuting the coordinates. The endomorphism ϕ_d of raising each coordinate to its d -th power commutes with this action, so it descends to a degree d^n endomorphism of $X := \mathbb{P}^n/G$.

According to the philosophy of [KL07], $\mathbb{P}^n/G$ should have terminal singularities for almost all G . The precise conditions are checked in Lemma 3. Some examples where $\mathbb{P}^n/G$ is not rational are given by [Sa85]. In his examples, G is a nonabelian group of order p^9 for some prime p and $G \subset S_{|G|+1}$ fixes the last coordinate and acts by the left G -action on itself on the first $|G|$ coordinates.

2. **Remark.** 1. These examples suggest the possibility that if a Fano variety X of Picard number 1 has terminal singularities and admits a degree > 1 endomorphism then X is a quotient of $\mathbb{P}^n$.

2. It seems likely that if G is large enough, then the ϕ_d are the only endomorphisms of $\mathbb{P}^n$ that commute with the G -action. Thus we may get examples of Fano varieties which have very few endomorphisms of degree > 1 .

3. **Lemma.** Let $G \subset S_n$ be a subgroup with the induced action on $\mathbb{P}^{n-1}$ by permuting the coordinates. Assume that none of the elements of G is conjugate (in S_n)

to one of the permutations $(12), (123)$ or $(12)(34)$. Then $\mathbb{P}^{n-1}/G$ has terminal singularities.

Proof. Let the order of g be m . Assume that the action of $g \in G$ on the set $\{1, 2, \dots, n\}$ gives cycles of length $n_1, n_2, \dots, n_k$ with $\sum_{i=1}^k n_i = n$. So $n_i | m$ and we write $m = n_i m_i$. After a linear transformation of the coordinates, g acts by

$$(t_1, t_2, \dots, t_n) \rightarrow (t_1, \xi^{m_1} t_2, \dots, \xi^{(n_1-1)m_1} t_{n_1}, \dots, \xi^{(n_k-1)m_k} t_n),$$

where ξ is a primitive m -th root of unity.

Now we compute the *age* of g at a fixed point x of g . (See [IR96], [Re02] for the definition and for the Reid-Tai criterion.) If a fixed point is on the chart $t_p \neq 0$, then, for each $1 \leq j \leq k$, the contribution of the cycle of length n_j to the age is the sum of all numbers c/m where $0 \leq c < m$ and c can be written as $c = km_j - p$ for some $k \in \mathbb{Z}$. This sum is bigger or equal to

$$\frac{m_j}{m}(0 + \dots + (n_j - 1)) = \frac{n_j - 1}{2}.$$

Therefore, $\text{age}_x(g) \geq \sum_{i=1}^k (n_i - 1)/2$ is always larger than 1 if g is not any one of the cases listed above. The Reid-Tai criterion implies that $\mathbb{P}^{n-1}/G$ has terminal singularities. $\square$

REFERENCES

- [Am97] Amerik, E.; Maps onto certain Fano threefolds. *Doc. Math.* **2** (1997), 195211
- [ARV99] Amerik, E.; Rovinsky, M.; Van de Ven, A.; A boundedness theorem for morphisms between threefolds. *Ann. Inst. Fourier (Grenoble)* **49** (1999), no. **2**, 405–415.
- [Ch08] Choe, I.; Endomorphisms of hypersurfaces of Fano manifolds of Picard number 1. to appear in *Mathematische Nachrichten*.
- [HM03] Hwang, J.; Mok, N.; Finite morphisms onto Fano manifolds of Picard number 1 which have rational curves with trivial normal bundles. *J. Algebraic Geom.* **12** (2003), no. **4**, 627–651.
- [HN08] Hwang, J.; Nakayama, N.; On endomorphisms of Fano manifolds of Picard number one. <http://www.kurims.kyoto-u.ac.jp/preprint/file/RIMS1628.pdf>
- [IR96] Ito, Y.; Reid, M.; The McKay correspondence for finite subgroups of $\text{SL}(3, \mathbb{C})$. *Higher-dimensional complex varieties (Trento, 1994)*, 221–240, *de Gruyter, Berlin*, 1996.
- [KL07] Kollár, J.; Larsen, M.; Quotients of Calabi-Yau varieties. arXiv:math/0701466. To appear in *Algebra, Arithmetic and Geometry - Manin Festschrift*
- [NZ07] N. Nakayama; D. Zhang; Polarized Endomorphisms of Complex Normal Varieties, Kyoto Univ. RIMS Preprint RIMS-1613. 2007
- [Re02] Reid, M. La correspondance de McKay. Séminaire Bourbaki, Vol. 1999/2000. *Astérisque* No. **276** (2002), 53–72.
- [Sa85] Saltman, D.; Noether’s problem over an algebraically closed field. *Invent. Math.* **77** (1984), no. **1**, 71–84.
- [Zh08] D. Zhang; Polarized endomorphisms of uniruled varieties. <http://www.math.nus.edu.sg/~matzdz/>